

IDENTIFYING ASH TREES

Take this guide to each tree on your property to identify ash



Ridged Bark:

On mature trees (left), bark is tight and displays patterns of diamond shaped ridges. On young trees (right), bark is relatively smooth.



Compound 'Opposite' Leaves:

Leaves contain 5 to 11 leaflets with smooth or toothed margins (tips). Leaflets are positioned opposite with one at the top.



Seeds:

When present, seeds usually hang in clusters and are dry and oar-shaped.



'Opposite' Branches:

Branches and buds are directly across from each other rather than staggered. However, due to the death and grooming of individual branches, it is possible that not every branch will be opposite.



WHAT IS THE EMERALD ASH BORER?

The Emerald Ash Borer is a metallic green wood-boring beetle of about 1 to 1.5 cm in length that attacks all native species of ash trees, typically killing them in 2 to 3 years. Its larva bore tunnels inside the tree, feeding off the inner bark until the tree dies.

Native to northeastern Asia, the pest was first discovered in Ontario in the Windsor area in 2002. Since then, infested ash trees have been discovered in Essex, Lambton, Elgin and Middlesex Counties, and in the Municipality of Chatham-Kent.

RECOGNIZING INFESTED ASH TREES

Infested ash trees often exhibit the following symptoms



Crown Dieback:

Severely attacked trees may exhibit crown dieback as the canopy dies from the top down. Leaves may wilt or turn yellow during the growing season.



Bark Cracks:

Vertical splits of 7 - 10 cm are often present over larval galleries. These are often more noticeable on young trees that do not already have splits from growth-related expansion.



Woodpeckers:

Woodpeckers feed on the larvae under the bark. Look for increased Woodpecker feedings or signs of their probing in the bark.



Exit Holes:

Once fully mature, the adult beetles emerge through exit holes they chew through the bark. These holes are distinctly D-shaped and are 3.5 to 4 mm across.



Tunnels:

Winding S-shaped larval tunnels snake under the bark where larvae bore channels. Removing the bark exposes larvae and sawdust-filled galleries.